

**Dictionnaire des maladies
éponymiques et des observations
princeps : oeuf : développement des
cellules nerveuses (embryologie)**

**HARRISON, Ross Granville. -
Observations on the living developing
nerve fiber.**

In : The Anatomical record., 1907, Vol. 1, pp. 116-18

OBSERVATIONS ON THE LIVING DEVELOPING NERVE
FIBER.¹

BY

ROSS G. HARRISON.

The immediate object of the following experiments was to obtain a method by which the end of a growing nerve could be brought under direct observation while alive, in order that a correct conception might be had regarding what takes place as the fiber extends during embryonic development from the nerve center out to the periphery.

The method employed was to isolate pieces of embryonic tissue, known to give rise to nerve fibers, as for example, the whole or fragments of the medullary tube, or ectoderm from the branchial region, and to observe their further development. The pieces were taken from frog embryos about 3 mm. long at which stage, *i. e.*, shortly after the closure of the medullary folds, there is no visible differentiation of the nerve elements. After carefully dissecting it out, the piece of tissue is removed by a fine pipette to a cover slip upon which is a drop of lymph freshly drawn from one of the lymph-sacs of an adult frog. The lymph clots very quickly, holding the tissue in a fixed position. The cover slip is then inverted over a hollow slide and the rim sealed with paraffine. When reasonable aseptic precautions are taken, tissues will live under these conditions for a week and in some cases specimens have been kept alive for nearly four weeks. Such specimens may be readily observed from day to day under highly magnifying powers.

While the cell aggregates, which make up the different organs and organ complexes of the embryo, do not undergo normal transformation in form, owing, no doubt, in part, to the abnormal conditions of mechanical tension to which they are subjected; nevertheless, the individual tissue elements do differentiate characteristically. Groups of epidermis cells round themselves off into little spheres or stretch out into long bands, their cilia remain active for a week or more and a typical cuticular border develops. Masses of cells taken from the myotomes differentiate into muscle fibers showing fibrillæ with typical striations. When portions of myotomes are left attached to a piece of the medullary cord the muscle fibers which develop will, after two or three days, exhibit frequent contractions. In pieces of nervous tissue numerous fibers are formed, though, owing to the fact that they are developed largely within the mass

¹Read before the Society for Experimental Biology and Medicine at the 23d meeting, New York, May 22, 1907.

of transplanted tissue itself, their mode of development cannot always be followed. However, in a large number of cases fibers were observed which left the mass of nerve tissue and extended out into the surrounding lymph-clot. It is these structures which concern us at the present time.

In the majority of cases the fibers were not observed until they had almost completed their development, having been found usually two, occasionally three, and once or twice four days after isolation of the tissue. They consist of an almost hyaline protoplasm, entirely devoid of the yolk granules, with which the cell-bodies are gorged. Within this protoplasm there is no definiteness of structure; though a faint fibrillation may sometimes be observed and faintly-defined granules are discernible. The fibers are about 1.5-3 μ thick and their contours show here and there irregular varicosities. The most remarkable feature of the fiber is its enlarged end, from which extend numerous fine simple or branched filaments. The end swelling bears a resemblance to certain rhizopods and close observation reveals a continual change in form, especially as regards the origin and branching of the filaments. In fact, the changes are so rapid that it is difficult to draw the details accurately. It is clear we have before us a mass of protoplasm undergoing amoeboid movements. If we examine sections of young normal embryos shortly after the first nerves have developed, we find exactly similar structures at the end of the developing nerve fibers. This is especially so in the case of the fibers which are connected with the giant cells described by Rohon and Beard.

Still more instructive are the cases in which the fiber is brought under observation before it has completed its growth. Then it is found that the end is very active and that its movement results in the drawing out and lengthening of the fiber to which it is attached. One fiber was observed to lengthen almost 20 μ in 25 minutes, another over 25 μ in 50 minutes. The longest fibers observed were 0.2 mm. in length.

When the placodal thickenings of the branchial region are isolated, similar fibers are formed and in several of these cases they have been seen to arise from individual cells. On the other hand, other tissues of the embryo, such as myotomes, yolk endoderm, notochord, and indifferent ectoderm from the abdominal region do not give rise to structures of this kind. There can, therefore, be no doubt that we are dealing with a specific characteristic of nervous tissue.

It has not as yet been found possible to make permanent specimens which show the isolated nerve fibers completely intact. The structures are so delicate that the mere immersion in the preserving fluid is sufficient

to cause violent tearing and this very frequently results in the tearing away of the tissue in its entirety from the clot. Nevertheless, sections have been cut of some of the specimens and nerves have been traced from the walls of the medullary tube but they were in all cases broken off short.

In view of this difficulty an effort, which resulted successfully, was made to obtain permanent specimens in a somewhat different way. A piece of medullary cord about four or five segments long was excised from an embryo and this was replaced by a cylindrical clot of proper length and caliber which was obtained by allowing blood or lymph of an adult frog to clot in a capillary tube. No difficulty was experienced in healing the clot into the embryo in proper position. After two, three, or four days the specimens were preserved and examined in serial sections. It was found that the funicular fibers form the brain and anterior part of the cord, consisting of naked axones without sheath cells, had grown for a considerable distance into the clot.

These observations show beyond question that the nerve fiber develops by the outflowing of protoplasm from the central cells. This protoplasm retains its amoeboid activity at its distal end, the result being that it is drawn out into a long thread which becomes the axis cylinder. No other cells or living structures take part in this process.

The development of the nerve fiber is thus brought about by means of one of the very primitive properties of living protoplasm, amoeboid movement, which, though probably common to some extent to all the cells of the embryo, is especially accentuated in the nerve cells at this period of development.

The possibility becomes apparent of applying the above method to the study of the influences which act upon a growing nerve. While at present it seems certain that the mere outgrowth of the fibers is largely independent of external stimuli, it is, of course, probable that in the body of the embryo there are many influences which guide the moving end and bring about contact with the proper end structure. The method here employed may be of value in analyzing these factors.

APPOINTMENTS.

Dr. Harold D. Senior, of the Wistar Institute of Anatomy, has accepted the position of Associate Professor of Anatomy at Syracuse University and will take up the duties of his new position in September. During the past winter Dr. Senior has been filling the chair of anatomy at Washington University, St. Louis, carrying on the work of Dr. R. J. Terry, who is at Harvard University on leave of absence.